

DATE: March 9, 2010

TERM TEST 2

TITLE PAGE

DEPARTMENT & COURSE NO: MATH 2300TIME: 75 minutesEXAMINATION: Linear Algebra IIEXAMINER: Borgersen

NAME: (Print in ink) _____

STUDENT NUMBER: _____

SIGNATURE: (in ink) _____

(I understand that cheating is a serious offense)

INSTRUCTIONS TO STUDENTS:

This is a 75 minute exam. **Please show your work clearly.**

No texts or notes are permitted. No calculators are permitted. Cell phones, electronic translators, and other electronic devices are **not** permitted.

This exam has a title page and 11 pages of questions, including 2 blank pages for rough work and 1 page showing the axioms of a vector space. Please check that you have all the pages. You may remove the blank pages and axiom page if you want, but be careful not to loosen the staple.

The value of each question is indicated beside the statement of the question. The total value of all questions is 78 points.

If you need more scrap paper, use the back of the question pages.

Question	Points	Score
1	10	
2	12	
3	10	
4	14	
5	4	
6	6	
7	10	
8	12	
Total:	78	

True or False Questions

1. [10 points] Are the following true or false? (Write "True" or "False" on the line to the right). **These are marked right minus wrong, so if you don't know, don't guess.** One mark each.

(a) Let $B = \{\mathbf{v}_1, \dots, \mathbf{v}_n\}$ be a basis for a vector space V , and let $T : V \rightarrow V$ be an linear operator. Then $T(B) = \{T(\mathbf{v}_1), \dots, T(\mathbf{v}_n)\}$ must be a basis for V as well.

(a) _____

(b) Let $T : \mathbb{R}^4 \rightarrow \mathbb{R}^3$ be defined by $T(a, b, c, d) = (a, b, c)$. Then T is an isomorphism.

(b) _____

(c) Let A be an invertible matrix. Then T_A must be 1-to-1.

(c) _____

(d) The vector spaces P_4 and \mathbb{R}^4 are isomorphic.

(d) _____

(e) Let $T_A : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ be the linear transformation that reflects vectors in the xz -plane, and then cuts their lengths in half. Then

$$A = \begin{bmatrix} \frac{1}{2} & 0 & 0 \\ 0 & \frac{-1}{2} & 0 \\ 0 & 0 & \frac{1}{2} \end{bmatrix}.$$

(e) _____

(f) Let $T : \mathbb{R}^2 \rightarrow \mathbb{R}$ be defined by $T(a, b) = \|(a, b)\|$. Then T is an onto linear transformation.

(f) _____

(g) Let $T : P_2 \rightarrow P_1$ be defined as $T(f(x)) = f'(x)$. Then T is a linear transformation.

(g) _____

(h) Let $T : V \rightarrow W$ be a one-to-one linear transformation. Then $T^{-1} : T(V) \rightarrow V$ is an isomorphism.

(h) _____

(i) Let $T : V \rightarrow W$ be a one-to-one linear transformation. Then $\text{nullity}(T^{-1}) = 0$.

(i) _____

(j) Let $S = \{\mathbf{s}_1, \dots, \mathbf{s}_n\}$ be a set of vectors in V and let T be a linear transformation $T : V \rightarrow W$ be such that $T(S) = \{T(\mathbf{s}_1), \dots, T(\mathbf{s}_n)\}$ is a linearly independent set in W . Then S must have been linearly independent.

(j) _____

DATE: March 9, 2010

TERM TEST 2

PAGE: 2 of 11

DEPARTMENT & COURSE NO: MATH 2300TIME: 75 minutesEXAMINATION: Linear Algebra IIEXAMINER: Borgersen

Long Answer Questions2. Let $T : P_2 \rightarrow P_1$ be defined by

$$T(a + bx + cx^2) = (a + b)x.$$

(a) [8 points] Show that T is a linear transformation.

DATE: March 9, 2010

TERM TEST 2

PAGE: 3 of 11

DEPARTMENT & COURSE NO: MATH 2300TIME: 75 minutesEXAMINATION: Linear Algebra IIEXAMINER: Borgersen

Reminder: $T : P_2 \rightarrow P_1$ is defined by

$$T(a + bx + cx^2) = (a + b)x.$$

(b) [4 points] Find a basis for the range of T .

3. [10 points] Let $T : P_2 \rightarrow M_{2,2}$ be defined by

$$T(f(x)) = \begin{bmatrix} f(1) & f(0) \\ f(-1) & -f(0) \end{bmatrix}.$$

Find a set that spans $\ker(T)$, and find $\text{nullity}(T)$.

DATE: March 9, 2010

TERM TEST 2

DEPARTMENT & COURSE NO: MATH 2300

PAGE: 5 of 11

EXAMINATION: Linear Algebra IITIME: 75 minutesEXAMINER: Borgersen

4. Let $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be a linear transformation such that

$$T(1, 0) = (1, 1) \quad \text{and} \quad T(0, 1) = (1, -1).$$

(a) [8 points] Find a general formula for $T(x, y)$.

(b) [2 points] What is $T(3, 5)$?

(c) [4 points] Find a general formula for $(T \circ T)(x, y)$.

5. Let $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be defined as $T(x, y) = (0, 4y + x)$

(a) [2 points] Is T one-to-one? Justify your answer.

(b) [2 points] Is T onto? Briefly justify your answer.

6. Let T_A be the linear transformation associated with the matrix

$$A = \begin{bmatrix} 1 & 4 & 5 & 0 & 9 \\ 3 & -2 & 1 & 0 & -1 \\ -1 & 0 & -1 & 0 & -1 \\ 2 & 3 & 5 & 1 & 8 \end{bmatrix}.$$

(a) [2 points] What is the domain of T_A ?

(b) [2 points] What is the codomain of T_A ?

(c) [2 points] Is T_A one-to-one?

DATE: March 9, 2010

TERM TEST 2

PAGE: 7 of 11

DEPARTMENT & COURSE NO: MATH 2300

TIME: 75 minutes

EXAMINATION: Linear Algebra II

EXAMINER: Borgersen

General Proofs

7. [10 points] Let $T : V \rightarrow W$ be a linear transformation. Prove that the $\ker(T)$ is a subspace of V .

DATE: March 9, 2010

TERM TEST 2

PAGE: 8 of 11

DEPARTMENT & COURSE NO: MATH 2300TIME: 75 minutesEXAMINATION: Linear Algebra IIEXAMINER: Borgersen

8. [12 points] Let $B = \{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n\}$ be a basis for a vector space V , and let $T : V \rightarrow W$ be a linear transformation. Show that if $T(\mathbf{v}_1) = T(\mathbf{v}_2) = \dots = T(\mathbf{v}_n) = \mathbf{0}_W$, then T is the zero transformation (that is, for every $\mathbf{v} \in V$, $T(\mathbf{v}) = \mathbf{0}_W$).

The Axioms of a Vector Space

A set V together with addition " \oplus " and scalar multiplication " \cdot " is a **Vector Space** if and only if each of the following axioms hold:

A1. for every $\mathbf{u}, \mathbf{v} \in V$, $\mathbf{u} \oplus \mathbf{v} \in V$,

A2. for every $\mathbf{u}, \mathbf{v} \in V$, $\mathbf{u} \oplus \mathbf{v} = \mathbf{v} \oplus \mathbf{u}$,

A3. for every $\mathbf{u}, \mathbf{v}, \mathbf{w} \in V$, $\mathbf{u} \oplus (\mathbf{v} \oplus \mathbf{w}) = (\mathbf{u} \oplus \mathbf{v}) \oplus \mathbf{w}$,

A4. there exists an element $\mathbf{0} \in V$ such that for every $\mathbf{u} \in V$, $\mathbf{0} \oplus \mathbf{u} = \mathbf{u}$,

A5. for every $\mathbf{u} \in V$, there exists a " $-\mathbf{u}$ " $\in V$ such that $\mathbf{u} \oplus (-\mathbf{u}) = \mathbf{0}$,

M1. for every $\mathbf{u} \in V$ and $k \in \mathbb{R}$, $k \cdot \mathbf{u} \in V$,

M2. for every $\mathbf{u}, \mathbf{v} \in V$ and $k \in \mathbb{R}$, $k \cdot (\mathbf{u} \oplus \mathbf{v}) = k \cdot \mathbf{u} \oplus k \cdot \mathbf{v}$,

M3. for every $\mathbf{u} \in V$, and $k, m \in \mathbb{R}$, $(k + m) \cdot \mathbf{u} = k \cdot \mathbf{u} \oplus m \cdot \mathbf{u}$,

M4. for every $\mathbf{u} \in V$, and $k, m \in \mathbb{R}$, $k \cdot (m \cdot \mathbf{u}) = (km) \cdot \mathbf{u}$, and

M5. for every $\mathbf{u} \in V$, $1 \cdot \mathbf{u} = \mathbf{u}$.

UNIVERSITY OF MANITOBA

DATE: March 9, 2010

TERM TEST 2

PAGE: 10 of 11

DEPARTMENT & COURSE NO: MATH 2300

TIME: 75 minutes

EXAMINATION: Linear Algebra II

EXAMINER: Borgersen

SCRAP PAPER

UNIVERSITY OF MANITOBA

DATE: March 9, 2010

TERM TEST 2

PAGE: 11 of 11

DEPARTMENT & COURSE NO: MATH 2300

TIME: 75 minutes

EXAMINATION: Linear Algebra II

EXAMINER: Borgersen

SCRAP PAPER